

The Effectiveness of LOUD Crowd® Group Therapy for People with Parkinson's Disease

References

- [1] Parkinson's Disease Foundation. (n.d.). *Statistics*. Retrieved September 17, 2020, from <https://www.parkinson.org/Understanding-Parkinsons/Statistics>
- [2] Freed, D. (2020). *Motor speech disorders: Diagnosis and treatment* (3rd ed.). Plural Publishing.
- [3] Parkinson Voice Project. (2020). *Our speech therapy program*. Retrieved on September 20, 2020 from <http://www.parkinsonvoiceproject.org>
- [4] Parkinson Voice Project. (2020). *Online Speak Out!® and LOUD Crowd® training course*. <http://www.parkinsonvoiceproject.org>
- [5] Fox, C. M., Ebersbach, G., Ramig, L. O., & Sapir, S. (2012). LSVT Loud and LSVT Big: behavioral treatment programs for speech and body movement in Parkinson's disease. *Parkinson's Disease*. <https://doi.org/10.1155/2012/391946>
- [6] Ramig, L. O., Sapir, S., Fox, C. M., & Countryman, S. (2001). Changes in vocal loudness following intensive voice treatment (LSVT) in individuals with Parkinson's disease: A comparison with untreated patients and normal age-matched controls. *Movement Disorders*, *16*(1), 79-83.
- [7] Watts, C. R. (2016). A retrospective study of long-term treatment outcomes for reduced vocal intensity in Hypokinetic Dysarthria. *BMC Ear, Nose and Throat Disorders*, *16*(2). doi: 10.1186/s12901-016-0022-8
- [8] Lam, J., & Tjaden, K. (2016). Clear speech variants: An acoustic study of Parkinson's disease. *Journal of Speech, Language, and Hearing Research*, *59*(4), 631–646.
- [9] Tjaden, K., Sussman, J. E., & Wilding, G. E. (2014). Impact of clear, loud, and slow speech on scaled intelligibility and speech severity in Parkinson's disease and multiple sclerosis. *Journal of Speech, Language, and Hearing Research*, *57*(3), 779–792.
- [10] Cannito, M., Suiter, D., Beverly, D., Chorna, L., Wolf, T., & Pfeiffer, R. (2012). Sentence intelligibility before and after voice treatment in speakers with idiopathic Parkinson's disease. *Journal of Voice*, *26*(2), 214-219.
- [11] Dromey, C., Ramig, L. O., & Johnson, A. B. (1995). Phonatory and articulatory changes associated with increased vocal intensity in Parkinson's disease: A case study. *Journal of Speech and Hearing Research*, *38*(4), 751-764.
- [12] El Sharkawi, A., Ramig, L. O., Logemann, J. A., Pauloski, B. R., Rademaker, A. W., Smith, Pawlas, A., Baum, S., & Werner, C. (2002). Swallowing and voice effects of Lee Silverman Voice Treatment (LSVT): A pilot study. *Journal of Neurol Neurosurg Psychiatry*, *72*(1), 31-36.

- [13] Levy, E., Moya-Gale, G., Chang, Y., Freeman, K., Forrest, K., Brin, M., & Ramig, L. (2020). The effects of intensive speech treatment on intelligibility in Parkinson's disease: A randomized controlled trial. *The Lancet's EClinical Medicine*, 24, 1-11.
- [14] Miles, A., Jardine, M., Johnston, F., deLisle, M., Friary, P., & Allen, J. (2017). Effect of Lee Silverman Voice Treatment (LSVT LOUD®) on swallowing and cough in Parkinson's disease: A pilot study. *Journal of the Neurological Sciences*, 383, 180-187.
- [15] Moya-Gale, G., Goudarzi, A., Bayes, A., McAuliffe, M., Bulte, B., & Levy, E. (2018). The effects of intensive speech treatment conversational intelligibility in Spanish speakers with Parkinson's disease. *American Journal of Speech-Language Pathology*, 27(1), 154-165.
- [16] Ramig, L. O., Fox, C. M., & Sapir, S. (2004). Parkinson's disease: Speech and voice disorders and their treatment with the Lee Silverman Voice Treatment. *Seminar in Speech and Language*, 25(2), 169-180.
- [17] Sapir, S., Ramig, L. O., & Fox, C. M. (2011). Intensive voice treatment in Parkinson's disease: Lee Silverman Voice Treatment. *Expert Review of Neurotherapeutics*, 11(6), 815-830.
- [18] Sapir, S., Spiellman, J. L., Ramig, L. O., Story, B. H., & Fox, C. M. (2007). Effects of intensive voice treatment (the Lee Silverman Voice Treatment [LSVT]) on vowel articulation in dysarthric individuals with idiopathic Parkinson's disease: Acoustic and perceptual findings. *Journal of Speech, Language, and Hearing Research*, 50(4), 899-912.
- [19] Sauvageau, V. M., Roy, J., Langlois, M., & Macoir, J. (2015). Impact of the LSVT on vowel articulation and coarticulation in Parkinson's disease. *Clinical Linguistics & Phonetics*, 29(6), 424-440.
- [20] Ramig L, Mead C, Scherer R, Horii, Y., Larson, K., & Kohler, D. (1988) Voice therapy and Parkinson's disease: A longitudinal study of efficacy [Paper presentation]. Clinical Dysarthria Conference, San Diego, CA.
- [21] Fox, C. M., Morrison, C. E., Ramig, L. O., & Sapir, S. (2002). Current perspectives on the Lee Silverman Voice Treatment (LSVT) for individuals with idiopathic Parkinson disease. *American Journal of Speech-Language Pathology*, 11(2), 111-123.
- [22] Nakayama, K., Yamamoto, T., Oda, C., Sato, M., Murakami, T., & Horiguchi, S. (2020). Effectiveness of Lee Silverman Voice Treatment LOUD on Japanese-speaking patients with Parkinson's disease. *Rehabilitation Research and Practice*, 2020(3), 1-7. <https://doi.org/10.1155/2020/6585264>
- [23] Ramig, L. O., Countryman, S., O'Brien, C., Hoehn, M., & Thompson, L. (1996). Intensive speech treatment for patients with Parkinson's disease: Short- and long-term comparison of two techniques. *Neurology*, 47(6), 1496-504.
- [24] Ramig, L., Halpern, A., Spielman, J., Fox, C., & Freeman, K. (2018). Speech treatment in Parkinson's disease: Randomized controlled trial (RCT). *Movement Disorders*, 33(11), 1777-1791.

- [25] Ramig, L. O., Sapir, S., Countryman, S., Pawlas, A. A., O'Brien, C., Hoehn, M., & Thompson, L. L. (2001). Intensive voice treatment for patients with Parkinson's disease: A 2-year follow up. *Journal of Neurol Neurosurg Psychiatry*, 71(4), 493-498.
- [26] Ramig, L. O., Sapir, S., Fox, C. M., & Countryman, S. (2001). Changes in vocal loudness following intensive voice treatment (LSVT) in individuals with Parkinson's disease: A comparison with untreated patients and normal age-matched controls. *Movement Disorders*, 16(1), 79-83.
- [27] Sapir, S., Ramig, L. O., & Fox, C. M. (2008). Speech and swallowing disorders in Parkinson disease. *Current Opinion Otolaryngology Head and Neck Surgery*, 16(3), 205-210.
- [28] Sapir, S., Ramig, L. O., & Fox, C. M. (2011). Intensive voice treatment in Parkinson's disease: Lee Silverman Voice Treatment. *Expert Review of Neurotherapeutics*, 11(6), 815-830.
- [29] Sapir, S., Ramig, L. O., Hoyt, P., Countryman, S., O'Brien, C., & Hoehn, M. (2002). Speech loudness and quality 12 months after intensive voice treatment (LSVT) for Parkinson's disease: A comparison with an alternative speech treatment. *Folia Phoniatrica Logopaedica*, 54(6), 296-303.
- [30] Sapir, S., Spiellman, J. L., Ramig, L. O., Story, B. H., & Fox, C. M. (2007). Effects of intensive voice treatment (the Lee Silverman Voice Treatment [LSVT]) on vowel articulation in dysarthric individuals with idiopathic Parkinson's disease: Acoustic and perceptual findings. *Journal of Speech, Language, and Hearing Research*, 50(4), 899-912.
- [31] Wight, S., & Miller, N. (2015). Lee Silverman Voice Treatment for people with Parkinson's: Audit of outcomes in a routine clinic. *International Journal of Language & Communication Disorders*, 50(2), 215-225.
- [32] Yorkston, K., Hakel, M., Beukelman, D., & Fager, S. (2007). Evidence for effectiveness of treatment of loudness, rate, or prosody in dysarthria: A systematic review. *Journal of Medical Speech Language Pathology*, 15(2), 11-36.
- [33] Spielman, J., Ramig, L. O., Mahler, L., Halpern, A., & Gavin, W. J. (2007). Effects of an extended version of the Lee Silverman Voice Treatment on voice and speech in Parkinson's disease. *American Journal of Speech Language Pathology*, 16(2), 95-107.
- [34] Baumann, A., Nebel, A., Granert, O., Giehl, K., Wolff, S., Schmidt, W., Baasch, C., Schmidt, G., Witt, K., Deuschl, G., Hartwigsen, G., Zeuner, K., & van Eimeren, T. (2018). Neural correlates of hypokinetic dysarthria and mechanisms of effective voice treatment in Parkinson disease. *Neurorehabilitation and Neural Repair*, 32(12), 1055-1066.
- [35] Sackley, C. M., Smith, C. H., Rick, C. E., Brady, M. C., Ives, N., Patel, S., Woolley, R., Dowling, F., Patel, R., Roberts, H., Jowett, S., Wheatley, K., Kelly, D., Sands, G., & Clarke, C. E. (2018). Silverman Voice Treatment versus standard speech and language therapy versus control in Parkinson's disease: A pilot randomized controlled trial. *Pilot and Feasibility Studies*, 4, Article 30. <https://doi.org/10.1186/s40814-017-0222-z>

- [36] Spielman, J., Mahler, L., Halpern, A., Gilley, P., Klepitskaya, O., & Pamig, L. (2011). Intensive voice treatment (LSVT LOUD) for Parkinson's disease following deep brain stimulation of the subthalamic nucleus. *Journal of Communication Disorders*, 44(6), 688-700.
- [37] Wohlert, A. (2004). Service delivery variables and outcomes of treatment for hypokinetic dysarthria in Parkinson disease. *Journal of Medical Speech-Language Pathology*, 12(4), 235-239.
- [38] Searl, J., Wilson, K., Haring, K., Dietsch, A., Lyons, K., & Pahwa, R. (2011). Feasibility of group voice therapy for individuals with Parkinson's disease. *Journal of Communication Disorders*, 44(2), 719-732.
- [39] Traverse, C. (2016). The impact of group format therapy on voice in Parkinson's disease: A pilot project. *Canadian Journal of Speech-Language Pathology and Audiology*, 20(1), 31-49.
- [40] LSVT Global, 2020. "Reinforcing LSVT Treatment Gains". Retrieved on October 14, 2020 from https://lsvtglobal.secure.force.com/portal/Therapists_Professionals
- [41] Behrman, A., Cody, J., Elandary, S., Flom, P., & Chitnis, S. (2020). The effect of SPEAK OUT! and The LOUD Crowd on dysarthria due to Parkinson's disease. *American Journal of Speech-Language Pathology*, 29(3), 1448-1465.
- [42] Levitt, J. S. (2014a). A case study: The effects of the "Speak Out!®" voice program for Parkinson's disease. *International Journal of Applied Science and Technology*, 4(2), 20-28.
- [43] Behrman, A., Wilson, M., Cody, J., Elandary, S., & Chitnis, S. (2019). *The Effect of Intentional Speech on Prosody in Parkinson Disease* [Poster presentation]. American Speech-Language-Hearing Association Convention, Orlando, FL.
- [44] Boutsen, F., Park, E., Dvorak, J., & Cid, C. (2018). Prosodic improvement in persons with Parkinson disease receiving SPEAK OUT!® voice therapy. *Folia Phoniatica Et Logopaedica*, 70(2), 51-58.
- [45] Cleveland, L., Pimentel, J., Talarico, M. & Solomon, J. (2015). *Vocal intensity and speech intelligibility pre- and post-SPEAK OUT!® treatment* [Poster presentation]. American Speech-Language-Hearing Association Convention, Denver, CO.
- [46] Levitt, J.S., Walker-Batson, D. (2018). The effects of the "Speak With Intent" instruction for individuals with Parkinson's disease. *Journal of Communication Disorders and Assistive Technology*, 2(1), 1-15.
- [47] Elman, R. J., & Bernstein-Ellis, E. (1999). The efficacy of group communication treatment in adults with chronic aphasia. *Journal of Speech, Language and Hearing Research*, 42(2), 411-419.
- [48] Levitt, J. S., Chitnis, S., & Walker-Batson, D. (2015). The effect of the "SPEAK OUT!®" voice program for Parkinson's disease. *International Journal of Health Sciences*, 3(2), 13-19.

- [49] Levitt, J. S. (2014b). *The effects of the “Speak Out!®” and “Loud Crowd®” voice programs for Parkinson’s disease* [Poster presentation]. American Speech, Language, and Hearing Association Convention, Orlando, FL.
- [50] Darling-White, M., & Huber, J. (2017). The impact of expiratory muscle strength training on speech breathing in individuals with Parkinson's disease: A preliminary study. *American Journal of Speech-Language Pathology*, 26(4), 1159-1166.
- [51] Reyes, A., Castillo, A., Castillo, J., Cornejo, I., & Cruickshank, T. (2019). The effects of respiratory muscle training on phonatory measures in individuals with Parkinson's disease. *Journal of Voice*. <https://doi.org/10.1016/j.jvoice.2019.05.001>
- [52] Haneishi, E. (2001). Effects of a music therapy voice protocol on speech intelligibility, vocal acoustic measures, and mood of individuals with Parkinson's disease. *Journal of Music Therapy*, 38(4), 273-290.
- [53] Higgins, A., & Richardson, K. (2019). The effects of a choral singing intervention on speech characteristics in individuals with Parkinson’s disease: An exploratory study. *Communication Disorders Quarterly*, 40(4), 195-205.
- [54] Tamplin, J., Morris, M., Marigliani, C., Baker, F., & Vogel, A. (2019). ParkinSong: A controlled trial of singing-based therapy for Parkinson’s disease. *Neurorehabilitation and Neural Repair*, 33(6), 453-463.
- [55] Yinger, O., & Lapointe, L. (2012). The effects of participation in a group music therapy voice protocol (G-MTVP) on the speech of individuals with Parkinson's disease. *Music Therapy Perspectives*, 30(1), 25-31.
- [56] Elefant, C., Baker, F., Lotan, M., Lagesen, S., & Skeie, G. (2012). The effect of group music therapy on mood, speech, and singing in individuals with Parkinson's disease — A feasibility study. *Journal of Music Therapy*, 49(3), 278-302.
- [57] Shih, L., Piel, J., Warren, A., Kraics, L., Silver, A., Vanderhorst, V., . . . Tarsy, D. (2012). Singing in groups for Parkinson's disease (SING-PD): A pilot study of group singing therapy for PD-related voice/speech disorders. *Parkinsonism and Related Disorders*, 18(5), 548-552.
- [58] Radad K, Gille G, Rausch W. (2005). Short review on dopamine agonists: Insight into clinical and research studies relevant to Parkinson's disease. *Pharmacological Reports*. 57, 701-712.
- [59] Ho, A., Bradshaw, J., & Iansek, R. (2008). For better or worse: The effect of levodopa on speech in Parkinson's disease. *Movement Disorders*, 23(4), 574-580.
- [60] De Letter, M., Santens, P., De Bodt, M., Van Maele, G., Van Borsel, J., & Boon, P. (2007). The effect of levodopa on respiration and word intelligibility in people with advanced Parkinson's disease. *Clinical Neurology and Neurosurgery*, 109(6), 495-500.
- [61] Plowman-Prine, E., Okun, M., Sapienza, C., Shrivastav, R., Fernandez, H., Foote, K., . . . Rosenbek, J. (2009). Perceptual characteristics of Parkinsonian speech: A comparison of the pharmacological effects of levodopa across speech and non-speech motor systems. *Neurorehabilitation*, 24(2), 131-144.

- [62] Fabbri, M., Guimarães, I., Cardoso, R., Coelho, M., Guedes, L., Rosa, M., . . . Ferreira, J. (2017). Speech and voice response to a levodopa challenge in late-stage Parkinson's disease. *Frontiers in Neurology*, 8, 432.
- [63] Skodda, S. (2012). Effect of Deep Brain Stimulation on Speech Performance in Parkinson's Disease. *Parkinson's Disease*, 10.
- [64] Krack, Paul, Batir, Alina, Blercom, Nadege van, Chabardes, Stephan, Fraix, Valerie, Ardouin, Claire, . . . Pollak, Pierre. (2003). Five-year follow-up of bilateral stimulation of the subthalamic nucleus in advanced Parkinson's disease. *The New England Journal of Medicine*, 349(20), 1925-19234.
- [65] Limousin, Patricia, Krack, Paul, Pollak, Pierre, Benazzouz, Abdelhamid, Ardouin, Claire, Hoffmann, Dominique, & Benabid, Alim-Louis. (1998). Electrical stimulation of the subthalamic nucleus in advanced Parkinson's disease. *The New England Journal of Medicine*, 339(16), 1105-1111.
- [66] Pinto, S., Gentil, M., Fraix, V., Benabid, A.L., Pollak, P. (2003) Bilateral subthalamic stimulation effects on oral force control in Parkinson's disease. *Journal of Neurology*, 250(2), 179–187.
- [67] Tripoliti, Zrinzo, Martinez-Torres, Frost, Pinto, Foltynie, . . . Limousin. (2011). Effects of subthalamic stimulation on speech of consecutive patients with Parkinson disease. *Neurology*, 76(1), 80-86.
- [68] Rouaud, T., Dondaine, T., Drapier, S., Haegelen, C., Lallement, F., Péron, J., . . . Vérin, M. (2010). Pallidal stimulation in advanced Parkinson's patients with contraindications for subthalamic stimulation. *Movement Disorders*, 25(12), 1839-1846.
- [69] Hariz, M.I., Rehncrona, S., Quinn, N.P., Speelman, J.D., Wensing, C. (2008) Multicenter study on deep brain stimulation in Parkinson's disease: an independent assessment of reported adverse events at 4 years. *Movement Disorders*, 23(3), 416–421.
- [70] Karlsson, F., Blomstedt, P., Olofsson, K., Linder, J., Nordh, E., Van Doorn, J. (2012) Control of phonatory onset and offset in Parkinson patients following deep brain stimulation of the subthalamic nucleus and caudal zona incerta. *Parkinsonism & Related Disorders*, 18, 824–827.
- [71] Lundgren, Saeys, Karlsson, Olofsson, Blomstedt, Linder, . . . Van Doorn. (2011). Deep Brain Stimulation of Caudal Zona Incerta and Subthalamic Nucleus in Patients with Parkinson's Disease: Effects on Voice Intensity. *Parkinson's Disease*, 2011(2011), 658956.
- [72] Kleiner-Fisman, G., Herzog, J., Fisman, D., Tamma, F., Lyons, K., Pahwa, R., . . . Rezai, Ali R. (2006). Subthalamic nucleus deep brain stimulation: Summary and meta-analysis of outcomes. *Movement Disorders*, 21(S14), S290-S304.